
CENTER FOR RECONSTRUCTIVE URETHRAL SURGERY



e-mail: info@urethralcenter.it

Websites: www.uretra.it www.urethralcenter.it



Nueva clasificación de las estenosis de uretra y algoritmo simplificado del tratamiento

Based-evidence Medicine

Metanalysis review

Guidelines

Consensus Conferences



Urological literature on urethral stricture

- ***** Few prospective, randomized studies
- **No homogeous series of patients**
- **❖** Small sample size − no statistical significance



Can Reconstructive Urethral Surgery Proceed Without Randomised Controlled Trials?

Guido Barbaglia, Massimo Lazzerib,*

What has history taught us? The history of urethral reconstruction is based on observational studies that appear today to be disarming in their apparent ease. In 1968, Orandi introduced the concept of penile skin flap urethroplasty, publishing a paper consisting merely of two pages and two black and white figures and that lacked any outcome assessment [10]. Oral mucosa has become the most popular substitute material in the treatment of urethral strictures, as it is readily available and easily harvested from the cheek, lip, or tongue, allowing for a concealed donor site scar and low oral morbidity [11-14]. Unfortunately, all the papers that have contributed to the widespread use of the oral mucosa graft are retrospective, not prospective, and they are not RCTs.

^{*}Center for Reconstructive Urethral Surgery, Arezzo, Italy

Department of Urology, Santa Chiara Hospital (GIOMI group), Florence, Italy

Personal opinion of high-volume surgeon from high-volume Center for Urethral Surgery (± 400 urethral surgeries for year)

Data published in peer review scientific Journals

Data fully available and updated in our websites every two years



Opposing Views

Urethral Strictures is Now an Open Surgical Disease

Allen Morey

Cost-Effective Approach to Short Bulbar Urethral Strictures Supports Single Internal Urethrotomy Before Urethroplasty

Hunter Wessels

J Urol 2009; 181: 953-955


New classification of anterior urethral strictures





Simple penile urethral stricture






- 1. The penis and genitalia are normal
- 2. Practicable urethral plate
- 3. No Lichen sclerosus
- 4. No failed hypospadis repair
- 5. No prior failed urethroplasty


Complex penile urethral stricture







- 1. Genital lichen sclerosus
- 2. Failed hypospadias
- 3. Prior failed urethroplasty
- 4. No practicable urethral plate








Wide urethral plate










































www.uretra.it
Websites: www.urethralcenter.it



Narrow urethral plate

















One-stage penile urethroplasty: graft or flap?

❖ NO sure data available from literature

❖ NO comparative studies including evaluation of post-operative sexual and aesthetic dysfuctions and patient satisfaction


Surgeon background and preference








Panurethral stricture
















Surgical repair of complex penile urethral stricture

Two-stage repair





First-stage - oral mucosa









39% of our patients required new surgical revision of the grafted area







Eur Urol 2006; 49: 887-895

Second stage - primary closure














First stage – Johanson







First stage – Johanson







Second stage - oral graft inlay





Second stage - oral graft inlay





Second stage - oral graft inlay














New classification of anterior urethral strictures



Etiology of bulbar urethral stricture




Catheter



Instrumentation



Infection





Trauma


Open or blunt perineal trauma












Urethral lesion requiring emergency treatment






- 1. Obliterative stricture
- 2. No practicable urethral plate
- 3. Scar tissue
- 4. Short or long fibrous gap






- 1. Etiology: idiopathic catheter instrumentation –infection other
- 2. Practicable urethral plate (3 Fr. Sensor guidewire)
- 3. No fibrous gap



Reconstruction of the bulbar urethra in traumatic stricture



Simple lithotomy position




Allen stirrups




Sequential inflatable compression sleeves

1 - 2 cm traumatic bulbar urethral stricture









www.uretra.it Websites: www.urethralcenter.it


End-to-end anastomosis






End-to-end anastomosis






2 - 4 cm traumatic bulbar urethral stricture








Augmented anastomotic repair






Augmented anastomotic repair





Augmented anastomotic repair







Reconstruction of the bulbar urethra in non-traumatic stricture






Insert Sensor 3 Fr. guidewire





Inject methylene blue inside the urethra

(G. Webster)






Calibrate the distal urethra and identify the distal stop

Non-traumatic bulbar urethral stricture Ventral or dorsal graft?



Distal


Dorsal onlay








Proximal


Ventral onlay





























Webster's technique






























































































Reconstruction of the bulbar urethra after previous failed urethroplasty




































Definitive perineal urethrostomy











Conclusions

In the era of robotic surgery, it is also time to change urethral surgery!

❖ Increase the use of minimally invasive techniques in urethroplasty, reducing the incidence of complications and improving patient quality of life.



Register now!

www.webon.uretra.it