

### FAILED HYPOSPADIAS REPAIR - one stage

type of repair	number	%	primary outcome			
			success	%	failure	%
circumcision	5	2,5%	5	100,0%		
removal of skin scrotal chordee	1	0,5%	1	100,0%		
circumcision and meatotomy	2	1,0%	2	100,0%		
meatoplasty	8	3,9%	4	50,0%	4	50,0%
meatotomy	48	23,6%	31	64,6%	17	35,4%
Barcat	3	1,5%	2	66,7%	1	33,3%
Byars	1	0,5%	1	100,0%		
meatotomy and Nesbit	3	1,5%	3	100,0%		
meatotomy and fistula closure	1	0,5%			1	100,0%
fistula closure	24	11,8%	17	70,8%	7	29,2%
excision of diverticulum	1	0,5%	1	100,0%		
Nesbit	4	2,0%	4	100,0%		
Nesbit and meatoplasty	1	0,5%	1	100,0%		
end-to-end anastomosis	7	3,5%	3	42,8%	4	57,2%
internal urethrotomy	9	4,4%	5	55,6%	4	44,4%
perineostomy	26	12,8%	22	84,6%	4	15,4%
one-stage repair using skin flap	12	5,9%	6	50,0%	6	50,0%
one-stage repair using skin graft	11	5,4%	6	54,5%	5	45,5%
one-stage repair using oral mucosal graft	36	17,7%	27	75,0%	9	25,0%
<b>TOTAL</b>	<b>203</b>	<b>100,0%</b>	<b>141</b>	<b>69,1%</b>	<b>62</b>	<b>30,9%</b>