

FAILED HYPOSPADIAS REPAIR - one stage						
type of repair	number	%	primary outcome			
			success	%	failure	%
circumcision	5	2,2%	5	100,0%		
removal of skin scrotal chordee	1	0,4%	1	100,0%		
circumcision and meatotomy	1	0,4%			1	100,0%
meatoplasty	9	4,0%	5	55,6%	4	44,4%
meatotomy	53	22,8%	36	68,0%	17	32,0%
Barcat	3	1,3%	2	66,7%	1	33,3%
Byars	1	0,4%	1	100,0%		
meatotomy and Nesbit	3	1,3%	3	100,0%		
meatotomy and fistula closure	1	0,4%			1	100,0%
fistula closure	27	11,6%	19	70,4%	8	29,6%
excision of diverticulum	1	0,4%	1	100,0%		
Nesbit	4	1,7%	4	100,0%		
end-to-end anastomosis	7	3,0%	3	42,9%	4	57,1%
internal urethrotomy	11	4,7%	6	54,5%	5	45,5%
perineostomy	34	14,7%	28	82,4%	6	17,6%
one-stage repair using skin flap	9	4,0%	3	33,3%	6	66,7%
one-stage repair using skin graft	11	4,7%	6	54,5%	5	45,5%
one-stage repair using oral mucosal graft	51	22,0%	33	64,7%	18	35,3%
TOTAL	232	100,0%	156	70,6%	76	29,4%